

Research Note

Occurrence of Multiple Mycotoxins in Various Types of Rice and Barley Samples in Thailand

PIYALUK SINPHITHAKKUL,^{1,2} AMNART POAPOLATHEP,¹ NARUMOL KLANGKAEW,¹ KANJANA IMSILP,¹
 ANTONIO F. LOGRIECO,³ ZHAOWEI ZHANG,⁴ AND SARANYA POAPOLATHEP^{1,2*}

¹Department of Pharmacology, Faculty of Veterinary Medicine and ²Center for Advanced Studies for Agriculture and Food, Institute for Advanced Studies, Kasetsart University, Bangkok 10900, Thailand; ³Institute of Sciences of Food Production, National Research Council, Bari, Italy; and ⁴Oil Crops Research Institute, Chinese Academy of Agricultural Sciences, Wuhan 430062, People's Republic of China

MS 18-456: Received 27 September 2018/Accepted 30 January 2019/Published Online 22 May 2019

ABSTRACT

The prevalence of mycotoxins is often increased by climatic conditions prevailing in tropical regions. Therefore, consumers in tropical countries such as Thailand have a higher risk of mycotoxin exposure. Existing reports have revealed mycotoxin contamination in rice. This study was conducted to determine the occurrence of multiple mycotoxins in barley and nine types of rice sold in Thailand and to assess consumer health risk. A total of 300 samples collected from various markets in Thailand were analyzed for the presence of 16 mycotoxins using a QuEChERS (quick, easy, cheap, effective, rugged, and safe) procedure and a triple quadrupole mass spectrometer equipped with an electrospray ionization source. Of the 300 samples, 124 (41.33%) were contaminated with at least one mycotoxin, and 38.71% of the mycotoxin-positive samples were simultaneously contaminated with more than one toxin. The incidence of mycotoxin contamination differed among the rice and barley samples. Beauvericin, diacetoxyscirpenol, zearalenone, and aflatoxins were the most frequently found mycotoxins. However, the concentrations of regulated mycotoxins were below the regulatory limits. The assessed mycotoxin exposure does not represent a health risk for Thai consumers because the estimated exposure concentrations were lower than the tolerable daily intake values established by the Joint FAO/WHO Expert Committee on Food Additives. However, our findings suggest that continued monitoring of mycotoxin contamination in rice and barley and concomitant risk assessments are warranted.

HIGHLIGHTS

- Contamination with multiple mycotoxins was found in rice and barley.
- BEA, DAS, ZEA, and aflatoxins were the mycotoxins most frequently found in samples.
- The assessed mycotoxin exposure does not represent a health risk for consumers.

Key words: Barley; Liquid chromatography–tandem mass spectrometry; Mycotoxins; QuEChERS; Rice

Mycotoxins are toxic substances produced by certain species of fungi. Contamination of agricultural commodities with mycotoxins during pre- and postharvest processing is a global problem that leads to serious risks to human and animal health. Environmental conditions, especially high humidity and temperature, can favor fungal proliferation, resulting in such contamination (3). Consumers in tropical countries such as Thailand encounter high risk of mycotoxin exposure. Mycotoxin contamination occurs in a wide variety of agricultural products, especially seeds and grains such as rice (*Oryza sativa*), which is one of the most important staple foods for the world's population. In Thailand, rice is the most important food crop and plays an important role in the economic system. Thailand is one of the world's most important rice producers, producing rice for both domestic consumption and export, and is ranked

among the top five Asian rice exporters (8). However, rice is relatively susceptible to fungal contamination, which can produce mycotoxins (17, 24, 25). Mycotoxins are very persistent in food and cannot be completely eliminated during food processing (18). Human health risks are usually associated with the direct consumption of food products contaminated with mycotoxins.

To date, several reports of mycotoxin contamination in rice from various countries have been published (2, 22, 26, 29, 32). Some studies have revealed the occurrence of aflatoxin B₁ (AFB₁) in Thailand (4, 11, 21, 28). Considering the health impact of mycotoxin consumption, data on contamination of rice by multiple mycotoxins are required to better estimate the potential problem. The main aim of this study was to evaluate barley and nine types of rice consumed in Thailand for the occurrence of various mycotoxins: AFB₁, aflatoxin B₂ (AFB₂), aflatoxin G₁ (AFG₁), aflatoxin G₂ (AFG₂), ochratoxin A (OTA), citrinin (CTN), nivalenol (NIV), deoxynivalenol (DON), diacetox-

* Author for correspondence. Tel: (662) 579-7537; Fax: (662) 579-7537; E-mail: fvetsys@ku.ac.th.

yscirpenol (DAS), T-2 toxin (T2), fumonisin B₁ (FB₁), fumonisin B₂ (FB₂), zearalenone (ZEA), sterigmatocystin (STG), beauvericin (BEA), and alternariol (ATL). These data were then used to assess consumer health risks.

MATERIALS AND METHODS

Toxins and chemicals. The analytical standards of mycotoxins AFB₁, AFB₂, AFG₁, AFG₂, OTA, CTN, NIV, DON, DAS, T2, FB₁, FB₂, ZEA, STG, and BEA were purchased from Romer Labs (Tulln, Austria), and the standard for ATL was from Sigma-Aldrich (Taufkirchen, Germany). Primary secondary amine (PSA) and octadecyl (C18) were purchased from Macherey-Nagel (Düren, Germany). Sodium chloride and magnesium sulfate anhydrous (MgSO₄) of analytical grade were purchased from AppliChem GmbH (Darmstadt, Germany). Other reagents and chemicals were of analytical grade.

Sample collection. A total of 300 samples were collected, 30 samples of each type of grain: black sticky rice, brown rice, GABA (γ -aminobutyric acid) rice, Japanese rice, jasmine rice, red rice, riceberry rice, white rice, white sticky rice, and barley. Samples were randomly collected from supermarkets and retail shops in Bangkok, Thailand, April to July 2017. Different brands were selected to obtain a representative sample of products sold in supermarkets in Bangkok. The samples were stored at 4°C until analyzed.

QuEChERS sample extraction and cleanup. The analysis was performed using a QuEChERS (quick, easy, cheap, effective, rugged, and safe) method modified from those of Arroyo-Manzanares et al. (1) and Jettanajit and Nhujak (11). A 1.0-g portion of a rice or barley sample was mixed with 5 mL of Milli-Q water and 5 mL of 10% (v/v) acetic acid in acetonitrile for 2 to 3 min with a vortex mixer. Then 2.0 g of MgSO₄, 0.5 g of NaCl, 0.5 g of sodium citrate tribasic dehydrate, and 0.25 g of sodium citrate dibasic sesquihydrate were added, and the mixture was shaken for 1 to 2 min. The acetonitrile fraction was separated by centrifugation at 1,968 \times g for 5 min, and 2 mL of the supernatant was transferred into a tube containing 300 mg of MgSO₄, 50 mg of C18, and 25 mg of PSA and shaken for 2 to 3 min. This mixture was separated by centrifugation at 1,968 \times g for 15 min, then 1 mL of supernatant was completely evaporated to dryness under a nitrogen stream at 40°C on a heating block. The residue was reconstituted with the mobile phase solution, injected onto a 0.22- μ m-pore-size syringe filter (Sartorius AG, Goettingen, Germany), and analyzed by liquid chromatography–tandem mass spectrometry (LC-MS/MS).

LC-MS/MS with electrospray ionization. A high-performance LC-MS/MS system with electrospray ionization was used to determine the presence of 16 mycotoxins in samples of barley and nine types of rice. The LC-MS/MS method was modified from that of Puangkhom et al. (23). The chromatographic separation was performed on a C18 column (50 by 4.6 mm, 1.8- μ m particle size; ZORBAX Eclipse plus RRHD, Agilent, Santa Clara, CA). The column was maintained at 40°C. The mobile phase consisted of 5 mM ammonium acetate with 0.2% acetic acid in water (mobile phase A) and 0.2% acetic acid in acetonitrile (mobile phase B) with a gradient elution protocol: 0 to 1 min, 10% B; 1 to 5 min, 10 to 95% B; 5 to 10 min, 95% B; and 10 to 12 min, 10% B. The column was reequilibrated for 3 min between injections. The mobile phase solution was filtered through a 0.22- μ m-pore-size membrane (Sartorius AG) and ultrasonically degassed prior to application. The flow rate was 0.3 mL/min, and the injection

volume was 5 μ L. The assay time was 14 min. The triple quadrupole mass spectrometer (6460 triple, Agilent, Waldbronn, Germany) was equipped with an electrospray ionization source run in both positive and negative ion modes under the multiple reaction monitoring mode, with two separate chromatographs analyzed per sample. The ionization source parameters were optimized as follows: capillary voltage, 3,500 V; gas temperature, 320°C; gas flow rate, 11 L/min; nebulizer, 50 lb/in². The parameters for the mass spectrometer were optimized as shown in Table 1.

Quantification and method validation. Validation of the LC-MS/MS method for AFB₁, AFB₂, AFG₁, AFG₂, OTA, CTN, NIV, DON, DAS, T2, FB₁, FB₂, ZEA, STG, BEA, and ATL was performed to assess the efficiency of this analytical method by investigating the recovery, repeatability, linear working range, limit of detection (LOD), limit of quantification (LOQ), accuracy, precision, and matrix effects in accordance with the guidelines of Commission Decision 2002/657/EC (5). We checked the suitability of the proposed method for the determination of mycotoxins in nine types of rice and in barley, and linearity, LOD, LOQ, precision, and trueness were determined using representative matrices from each category. The LOD and LOQ of the method were evaluated at signal/noise values of 3:1 and 10:1, respectively. Linear regression analysis was conducted for the 16 mycotoxin standards under the optimized LC-MS/MS conditions. Within-day recovery and precision (repeatability, expressed as percent relative standard deviation [RSD]) were determined by analyzing seven replicates of spiked samples containing the 16 mycotoxins at three quality control concentrations. The interday precision was determined by analyzing quality control samples on 5 days (one batch per day). The matrix-matched calibration curves were prepared in three replicates by spiking the working standard solution into blank samples to yield final concentrations of 1 to 50 μ g/kg (1, 2.5, 5, 10, 25, and 50 μ g/kg) for AFB₁, AFB₂, AFG₁, AFG₂, OTA, BEA, and STG; 1.5 to 100 μ g/kg (1.5, 5, 10, 25, 50, and 100 μ g/kg) for FB₁ and FB₂; 2 to 200 μ g/kg (2, 5, 10, 50, 100, and 200 μ g/kg) for ALT, CIT, and T2; 10 to 500 μ g/kg (10, 20, 50, 75, 100, 250, and 500 μ g/kg) for DAS, ZEA, and DON; and 100 to 1,000 μ g/kg (100, 200, 400, 500, 750, and 1,000 μ g/kg) for NIV. The matrix effects were determined by comparing the slope of six concentration calibration curves of the target analyte in postextraction spiked samples with those obtained in solvent alone.

Health risk evaluation. The type of mycotoxin exposure through rice consumption and data on the distribution of exposure concentrations within the studied population were combined with the average concentration for each contaminant under investigation. The mean and 97.5 percentile exposure (p97.5) values were evaluated, and a risk assessment was performed for average and high exposure. To compare dietary exposure concentrations with toxicological reference concentrations, which are expressed per kilogram of body weight, an average body weight of 60 kg was used (31). The estimate dietary exposure (EDI; μ g/kg of body weight per day) of the chemical residues were calculated with equation 1:

$$EDI = \sum RL_i \times \frac{A_i}{BW} \quad (1)$$

where RL_i is the mycotoxin residue in the rice or barley (μ g/kg), A_i is the rice or barley consumption rate (g per person per day), and BW is body weight (kg). Risk assessment is calculated by comparing the concentrations of residues detected with the provisional maximum tolerable daily intake (PMTDI). In this

TABLE 1. *MS/MS parameters for the determination of 16 mycotoxins*

Analyte	Precursor ion (<i>m/z</i>)	Product ions (<i>m/z</i>) ^a	Mean ion ratio ^b	Collision energy (eV)	Fragmentor (V)	Retention time (min)	Polarity
AFB ₁	313.07	285.1 241.0	0.75	21 35	150 150	6.32	Positive
AFB ₂	315.09	287.1 259.0	0.79	25 29	160 160	6.18	Positive
AFG ₁	329.07	311.0 243.0	0.91	25 43	160 160	6.16	Positive
AFG ₂	331.08	313.0 245.0	0.86	25 29	180 180	6.02	Positive
OTA	404.00	192.9 102.1	0.68	48 80	130 130	6.79	Positive
ZEA	319.16	283.0 187.0	0.83	5 17	80 80	7.24	Positive
BEA	801.40	784.3 244.1	0.52	13 35	160 160	9.13	Positive
FB ₁	722.40	352.5 334.4	0.89	40 45	160 160	5.59	Positive
FB ₂	706.30	336.2 318.3	0.75	35 40	200 200	5.99	Positive
T2	489.40	387.3 245.2	0.88	20 26	170 170	7.00	Positive
DAS	384.20	307.1 199.0	0.59	5 13	60 60	6.28	Positive
STER	325.10	309.9 281.0	0.68	25 35	160 160	7.46	Positive
ALT	259.10	187.9 160.1	0.81	25 33	240 240	6.08	Positive
CIT	251.10	233.1 205.1	0.21	3 20	120 120	5.79	Positive
DON	355.10	265.1 59.1	0.58	4 10	90 90	3.71	Negative
NIV	371.10	281.0 59.1	0.84	4 10	80 80	2.26	Negative

^a Bottom number of each couplet is the qualitative product ion.

^b Ratio of peak area of quantifier to qualifier transition.

TABLE 2. *Performance characteristics of the proposed method*

Analyte	White rice types			Colored rice types			Barley		
	LOD (µg/kg)	LOQ (µg/kg)	R ²	LOD (µg/kg)	LOQ (µg/kg)	R ²	LOD (µg/kg)	LOQ (µg/kg)	R ²
AFB ₁	0.28	0.92	0.9953	0.31	1.0	0.9974	0.25	0.85	0.9987
AFB ₂	0.21	0.71	0.9972	0.27	0.86	0.9979	0.21	0.63	0.9985
AFG ₁	0.14	0.47	0.9981	0.20	0.67	0.9985	0.16	0.52	0.9991
AFG ₂	0.17	0.52	0.9955	0.22	0.74	0.9962	0.17	0.55	0.9967
OTA	0.20	0.65	0.9991	0.20	0.69	0.9989	0.19	0.62	0.9991
ZEA	6.05	20.15	0.9967	10.6	35.2	0.9955	7.73	25.5	0.9973
BEA	0.20	0.68	0.9995	0.26	0.88	0.9989	0.15	0.51	0.9967
FB ₁	0.31	1.04	0.9994	0.47	1.55	0.9992	0.38	1.25	0.9959
FB ₂	0.44	1.48	0.9985	0.66	2.20	0.9987	0.53	1.77	0.9963
T2	0.76	2.53	0.9966	1.35	4.60	0.9936	0.98	3.26	0.9971
DAS	1.87	6.22	0.9985	2.63	8.75	0.9972	2.05	6.84	0.9989
STER	0.16	0.53	0.9978	0.22	0.75	0.9934	0.19	0.62	0.9959
DON	7.53	25.11	0.9956	10.3	34.5	0.9932	8.47	28.2	0.9948
NIV	26.80	89.34	0.9943	28.2	92.8	0.9935	25.6	85.3	0.9944
ALT	0.55	1.85	0.9989	0.68	2.30	0.9991	0.56	1.88	0.9981
CIT	0.64	2.14	0.9986	0.82	2.65	0.9985	0.67	2.23	0.9976

TABLE 3. Accuracy and precision in white rice types, colored rice types, and barley^a

Mycotoxin	White rice types									Colored rice types									Barley								
	Recovery (%) (n = 7)			Intraday precision (%RSD) (n = 7)			Interday precision (%RSD) (n = 18)			Recovery (%) (n = 7)			Intraday precision (%RSD) (n = 7)			Interday precision (%RSD) (n = 18)			Recovery (%) (n = 7)			Intraday precision (%RSD) (n = 7)			Interday precision (%RSD) (n = 18)		
	L1	L2	L3	L1	L2	L3	L1	L2	L3	L1	L2	L3	L1	L2	L3	L1	L2	L3	L1	L2	L3	L1	L2	L3	L1	L2	L3
AFB ₁	88.5	95.3	97.7	5.1	2.6	3.7	3.9	3.4	2.6	80.3	83.8	85.6	7.9	5.3	4.7	8.4	9.1	6.7	83.3	86.8	89.5	5.7	5.5	4.4	6.3	5.9	5.2
AFB ₂	87.6	95.9	95.7	2.5	3.1	1.9	2.1	1.4	1.4	81.2	86.7	87.9	6.9	4.8	3.1	7.8	7.9	7.6	86.4	90.1	88.6	4.5	4.9	4.6	5.9	5.4	4.9
AFG ₁	90.7	97.7	101	4.3	2.7	2.1	2.7	2.7	2.1	83.6	87.6	90.5	6.3	5.5	4.6	8.2	8.7	9.1	85.2	92.5	93.4	4.4	5.2	5.0	6.1	5.8	7.4
AFG ₂	85.7	93.5	94.3	3.4	2.7	1.7	1.8	2.2	1.3	80.7	83.5	88.6	7.4	4.7	3.5	8.5	9.3	7.2	88.1	85.7	90.3	5.9	6.1	4.7	7.5	7.1	7.7
OTA	92.9	94.1	99.5	2.6	2.1	3.2	2.4	1.9	2.5	84.5	87.7	88.3	5.8	6.1	4.7	7.4	8.5	7.2	92.8	90.4	91.7	4.3	3.8	2.9	4.6	4.2	5.6
ZEA	89.2	90.6	90.5	3.3	4.6	3.4	3.4	2.8	2.2	78.4	80.2	83.4	8.7	8.3	6.2	9.1	9.3	8.4	83.4	88.3	87.9	6.4	6.7	7.5	8.8	7.5	9.3
BEA	97.7	99.4	103	3.5	2.3	1.8	2.8	1.7	2.2	91.1	95.2	93.7	4.6	3.9	4.2	6.5	5.8	4.9	89.7	91.8	93.4	4.1	5.3	3.6	4.5	5.0	4.8
FB ₁	93.5	94.8	93.4	2.1	1.6	1.9	2.3	3.8	2.6	82.3	83.1	87.6	5.5	5.8	4.7	8.6	7.7	8.1	79.5	83.7	85.6	4.9	4.5	5.1	6.5	5.9	5.4
FB ₂	90.6	92.5	94.2	2.3	1.8	2.1	2.4	2.0	2.5	81.6	80.8	85.2	6.2	5.4	4.8	7.8	8.3	8.6	81.4	86.6	80.2	5.1	5.3	4.8	7.4	6.6	7.9
T2	92.0	93.2	102	4.2	3.3	4.2	3.0	2.4	3.7	85.3	84.4	87.9	5.7	4.9	3.9	9.7	9.6	7.3	81.3	84.8	88.6	3.8	4.4	3.9	4.2	3.2	5.1
DAS	95.4	96.6	98.6	2.7	2.7	2.0	2.5	1.7	1.8	87.7	89.6	86.8	4.5	4.7	4.2	6.9	5.7	5.5	85.7	87.9	90.4	4.2	3.9	4.5	4.3	4.5	3.8
STER	92.3	95.7	98.6	3.4	2.9	2.2	4.3	1.5	2.3	82.6	83.3	88.1	5.1	3.9	3.6	6.7	5.9	4.6	88.3	90.8	92.7	4.3	4.0	4.6	4.7	5.5	6.4
DON	81.6	88.1	89.4	6.3	5.7	4.9	5.6	4.9	6.4	73.6	72.2	78.6	8.3	6.5	5.7	9.1	7.7	7.2	78.2	82.6	80.3	7.8	6.5	7.9	7.5	8.1	7.6
NIV	79.5	83.8	82.1	7.7	5.4	4.9	4.4	5.4	3.2	75.2	78.3	79.8	8.5	7.6	6.1	9.8	8.1	7.0	74.5	79.7	83.6	6.9	7.1	8.3	8.1	8.5	7.7
ALT	89.0	91.1	95.4	3.0	2.2	2.7	2.6	1.9	2.2	83.2	85.9	86.2	4.2	3.8	3.5	6.7	6.0	5.9	86.8	89.4	92.4	4.6	4.8	3.9	5.3	6.2	6.5
CIT	90.2	92.2	93.7	3.5	1.7	2.3	2.4	2.1	1.7	82.5	85.7	80.4	4.7	4.5	3.6	6.6	7.2	5.5	85.3	88.8	90.6	4.7	3.2	2.8	5.7	5.1	3.2

^a Spiking levels. L1 for AFB₁, AFB₂, AFG₁, AFG₂, OTA, BEA, and STER: 1 µg/kg; for FB₁, FB₂, and ALT: 2 µg/kg; for T2 and CIT: 5 µg/kg; for DAS: 10 µg/kg; for ZEA and DON: 50 µg/kg; for NIV: 100 µg/kg. L2 for AFB₁, AFB₂, AFG₁, AFG₂, OTA, BEA, and STER: 10 µg/kg; for FB₁, FB₂, and ALT: 20 µg/kg; for T2 and CIT: 50 µg/kg; for DAS: 100 µg/kg; for ZEA and DON: 250 µg/kg; for NIV: 500 µg/kg. L3 for AFB₁, AFB₂, AFG₁, AFG₂, OTA, BEA, and STER: 40 µg/kg; for FB₁, FB₂, and ALT: 100 µg/kg; for T2 and CIT: 200 µg/kg; for DAS: 250 µg/kg; for ZEA and DON: 500 µg/kg; for NIV: 1,000 µg/kg.

TABLE 4. Matrix effects for each analyte in difference rice sample types and barley

Mycotoxin	Matrix effect (%)									
	White rice	Jasmine rice	Japanese rice	White sticky rice	Brown rice	GABA brown rice	Black sticky rice	Red rice	Riceberry rice	Barley
AFB ₁	87.27	86.72	86.41	83.35	79.32	81.37	78.16	80.15	78.84	86.48
AFB ₂	85.33	84.53	85.62	82.49	77.15	79.94	75.32	79.54	75.32	85.63
AFG ₁	83.79	84.65	85.35	84.11	80.26	80.65	79.47	78.36	78.52	88.32
AFG ₂	90.41	92.14	91.18	89.65	83.65	84.68	84.88	83.84	80.36	87.15
OTA	92.77	95.43	91.54	89.32	106.37	103.52	98.95	96.38	107.73	102.36
ZEA	68.32	70.46	72.36	71.64	62.95	65.71	64.32	63.11	64.12	70.17
BEA	94.54	98.59	95.15	90.52	111.32	107.58	109.68	107.22	101.35	101.44
FB ₁	91.71	93.45	93.16	90.36	78.65	76.23	73.42	79.33	72.59	101.48
FB ₂	84.23	84.18	82.47	81.51	73.77	75.85	72.53	76.16	72.65	106.38
T2	73.36	77.36	78.54	74.82	70.48	72.67	72.27	74.58	73.79	80.14
DAS	84.52	85.61	88.63	85.67	79.85	75.23	80.69	78.63	78.37	85.36
STER	86.78	88.79	87.49	82.36	82.63	80.85	81.37	82.37	80.16	87.83
DON	73.37	72.14	70.32	71.59	69.17	68.64	66.34	68.15	68.63	74.47
NIV	79.48	78.37	76.45	75.48	71.43	72.33	70.75	70.66	70.74	77.95
ALT	89.56	90.44	90.59	88.23	83.72	80.79	81.46	80.35	79.45	86.34
CIT	87.45	90.65	88.47	85.62	81.44	83.47	82.97	81.74	83.75	87.78

study, the hazard quotient (HQ) was used to indicate the long-term risk assessment. HQ can be calculated with equation 2:

$$HQ = \frac{EDI}{PMTDI} \times 100\% \quad (2)$$

When the value of HQ is >100%, the rice or barley involved should be considered a risk to consumers; when the value is <100%, the rice or barley is acceptable or low risk.

Because of the genotoxic and carcinogenic effects of aflatoxins, no TDI can be set. Therefore, the risk assessment calculation was performed separately by determination of the population at risk (cases of cancer per year per 100,000 persons) for primary liver cancer (12). The hepatitis B antigen (HBsAg)-positive prevalence rate was assumed to be 5% for the Thai population:

$$\text{Population risk} = \text{exposure} \times \text{average potency}$$

$$\begin{aligned} \text{Average potency} &= (0.3 \times \text{HBsAg}^+ \text{ rate}) \\ &\quad + (0.01 \times \text{HBsAg}^- \text{ rate}) \\ &= (0.3 \times 0.05) + (0.01 \times 0.95) \\ &= 0.0245 \text{ cases/year/100,000 persons} \end{aligned}$$

RESULTS

Optimization of LC-MS/MS. The optimization procedure was modified from that of Puangkham et al. (23). In both ionization modes, the highest sensitivity for all 16 mycotoxins was achieved using 5 mM ammonium acetate with 0.2% acetic acid in water (A: positive ion mode) and 5 mM ammonium acetate in 9:1 water-methanol (A: negative ion mode) with a mobile phase of 0.2% acetic acid in acetonitrile (positive ion mode) or methanol (negative ion mode). The MS parameters are shown in Table 1.

Method validation and quality assurance. The LC-MS/MS method used in this study had an adequate linearity range and adequate intra- and interday precision and accuracy for quantification of the 16 mycotoxins. Linearity

of the calibration curves for groups of white rice, groups of colored rice, and barley matrices, expressed as the determination coefficients (R^2), gave values that were all ≥ 0.9932 . The LOD and LOQ for mycotoxin detection in rice and barley were 0.14 to 26.80 and 0.47 to 89.34 $\mu\text{g}/\text{kg}$, respectively (Table 2). The mean recovery and intra- and interday precision (%RSD) of the spiked rice and barley samples are shown in Table 3. The percentage of extraction recovery of FB₁ reported in this study was higher than that reported by others (15). The differences in recovery rate may be associated with the smaller sample size used in our study or the use of alumina-neutral as a sorbent for the clean-up step in the previous study, which causes significant losses of acidic analytes. Considering matrix effects, we found that all compounds were subjected to ion suppression in white rice matrices. Both matrix enhancement and suppression were observed for target mycotoxin analytes in colored rice matrices and barley (Table 4). All compounds were subjected to ion suppression.

Occurrence of mycotoxins in rice and barley. The method described here was applied to a monitoring survey of 16 mycotoxins in barley and nine kinds of rice (black sticky rice, brown rice, GABA rice, Japanese rice, jasmine rice, red rice, riceberry rice, white rice, and white sticky rice). The summary of investigated compounds found in each variety is shown in Table 5. Overall, 124 (41.33%) of the total 300 samples were contaminated with at least one mycotoxin, and 38.71% of the mycotoxin-positive samples were simultaneously contaminated with more than one toxin. The incidence of mycotoxin contamination differed among the types of samples. BEA, DAS, ZEA, and aflatoxins were the mycotoxins most frequently found in all contaminated samples. AFB₁ was quantifiable in 20% (6 of 30) of black sticky rice samples, 10% (3 of 30) of riceberry rice samples, and 6.67% (2 of 30) of GABA brown rice samples; and AFG₁ was quantifiable in 3.33% (1

TABLE 5. Continued

Mycotoxin	Occurrence ^a	Type of sample									
		White rice	Jasmine rice	Japanese rice	White sticky rice	Brown rice	GABA brown rice	Black sticky rice	Red rice	Riceberry rice	Barley
NIV	No. (%) positive samples	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
	Mean concn (µg/kg)	0-26.8	0-26.8	0-26.8	0-26.8	0-26.8	0-26.8	0-26.8	0-26.8	0-26.8	0-26.8
ALT	No. (%) positive samples	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
	Mean concn (µg/kg)	0-0.6	0-0.6	0-0.6	0-0.6	0-0.6	0-0.6	0-0.6	0-0.6	0-0.6	0-0.6
CIT	No. (%) positive samples	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
	Mean concn (µg/kg)	0-0.6	0-0.6	0-0.6	0-0.6	0-0.6	0-0.6	0-0.6	0-0.6	0-0.6	0-0.6

^a Number of positive samples from a total of 30 samples of each type of rice or barley. Mean concentrations were calculated from the positive samples. LB, lower bound estimate calculated as the mean concentration for all samples; the undetected values and the unquantified values were replaced by zero and the LOD, respectively. UB, upper bound estimate calculated as the mean concentration for all samples; the undetected values and the unquantified values were replaced by the LOD and the LOQ, respectively.

of 30) of jasmine rice and white sticky rice samples. ZEA was detectable in 46.67% (14 of 30) of riceberry rice samples, 36.67% (11 of 30) of white sticky rice samples, 13.33% (4 of 30) of GABA brown rice and black sticky rice samples, 10% (3 of 30) of white rice samples, and 3.33% (1 of 30) of jasmine rice and Japanese rice samples. BEA was found in 66.67% (20 of 30) of black sticky rice samples, 60% (18 of 30) of GABA brown rice samples, 50% (15 of 30) of brown rice and riceberry rice samples, 46.67% (14 of 30) of red rice samples, 23% (7 of 30) of Japanese rice samples, 20% (6 of 30) of white rice and barley samples, and 16% (5 of 30) of jasmine rice and white sticky rice samples. DAS was found in 16.67% (5 of 30) of riceberry samples, 13.33% (4 of 30) of black sticky rice samples, 10% (3 of 30) of Japanese rice and red rice samples, 6.67% (2 of 30) of GABA brown rice samples, and 3.3% (1 of 30) of white rice samples. STER was detectable in 23.33% (7 of 30) of black sticky rice samples and 3.3% (1 of 30) of jasmine rice and riceberry rice samples. The mean concentrations and lower and upper estimates mycotoxin exposure from these grains are shown in Table 5.

The EDI of each residue of ZEA, DAS, STER, and BEA in rice and barley samples was calculated by multiplying either the mean or p97.5 of the consumption rate. The mean consumption rates for white rice, colored rice, and sticky rice were 255.66, 118.04, and 75.22 g per person per day, respectively, and the p97.5 values of the consumption rate were 720, 180, and 444 g per person per day, respectively (9, 19). The PMTDI value for ZEA was established by the Joint FAO/WHO Expert Committee on Food Additives (JECFA) for humans at 0.5 µg/kg of body weight (13). The HQ_{mean} and HQ_{97.5} values for ZEA, BEA, DAS, and STER in contaminated rice were all <100%. The EDI and HQ values for the contaminated rice and barley samples are shown in Table 6.

DISCUSSION

In our evaluation of mycotoxin contamination in rice and barley, the types of mycotoxins present were consistent with those reported in previous publications (14, 27). However, the mean concentration of AFB₁ was lower than those previously reported. AFB₁ has been analyzed in white rice, brown rice, basmati rice, whole grain rice, long grain rice, short grain rice, and puffed rice from Australia. Iqbal et al. (10) found that AFB₁ was quantifiable in 41% (14 of 34) and 54% (15 of 28) of white rice and brown rice samples, respectively. Nguyen et al. (20) reported that AFB₁ was detectable in 51% of positive samples from Vietnam, with a mean concentration of 3.31 µg/kg. Toteja et al. (30) reported high concentrations of aflatoxins (60 to 600 µg/kg) in parboiled rice samples. In our study, AFB₁ was quantifiable in GABA brown rice, black sticky rice, and riceberry rice but not in white rice and barley samples. This difference may associated with the milling and polishing process, which may reduce the AFB₁ concentration in white rice and barley. Our results are similar to those of Lim et al. (16), who reported low AFB₁ concentrations and no evidence of AFB₂, AFG₂, FB₁, FB₂, DON, T2, and OTA in rice samples collected from Thailand. In our study, ZEA was quantifiable

TABLE 6. Estimated dietary exposure and hazard quotient (HQ) values for mycotoxins through the consumption of rice and barley in Thailand

Mycotoxin	Mycotoxin-positive sample	Estimated dietary exposure ($\mu\text{g}/\text{kg}$ of BW/day)				EU maximum regulatory limit ($\mu\text{g}/\text{kg}$)	TDI or PMTDI ($\mu\text{g}/\text{kg}$ of BW/day)	HQ _{mean} (%)	HQ _{p97.5} (%)
		EDI _{mean}		EDI _{p97.5}					
		LB estimate ^a	UB estimate ^b	LB estimate	UB estimate				
ZEA	White rice	0.03	0.05	0.09	0.15	100	0.5	1.68–10.52	17.99–29.61
	Jasmine rice	0.01	0.03	0.02	0.09			1.58–6.18	4.46–17.42
	Japanese rice	0.01	0.05	0.02	0.12			1.66–6.13	4.32–17.28
	White sticky rice	0.04	0.04	0.23	0.26			7.87–8.76	46.48–51.72
	GABA rice	0.02	0.03	0.05	0.08			3.29–5.19	10.03–15.84
	Black sticky rice	0.01	0.01	0.05	0.09			1.67–2.88	9.86–17.02
	Riceberry rice	0.06	0.07	0.19	0.20			12.19–13.36	37.18–40.76
BEA	White rice	0.001	0.002	0.003	0.01	NA ^c	0.1 (TDI) ^d	1.17–1.80	3.29–5.06
	Jasmine rice	0.001	0.001	0.002	0.004			0.79–1.42	2.24–4.01
	Japanese rice	0.001	0.002	0.003	0.005			1.16–1.77	3.28–4.98
	White sticky rice	0.0003	0.0005	0.002	0.003			0.30–0.49	1.77–2.90
	Brown rice	0.002	0.002	0.007	0.007			2.22–2.40	6.78–7.34
	GABA rice	0.002	0.002	0.006	0.006			1.89–2.03	5.75–6.20
	Black sticky rice	0.001	0.001	0.01	0.018			1.41–1.49	8.34–8.80
	Red rice	0.002	0.002	0.005	0.006			1.72–1.91	5.25–5.84
	Riceberry rice	0.002	0.002	0.007	0.007			2.17–2.36	6.64–7.20
	Barley	0.001	0.001	0.004	0.004			0.90–1.20	2.76–6.65
DAS	White rice	0.001	0.01	0.002	0.028	NA	2	0.03–0.39	0.09–1.09
	Japanese rice	0.003	0.01	0.01	0.03			0.14–0.47	0.40–1.33
	GABA rice	0.001	0.005	0.004	0.01			0.07–0.23	0.21–0.70
	Black sticky rice	0.002	0.004	0.01	0.02			0.09–0.18	0.52–1.07
	Red rice	0.001	0.004	0.004	0.01			0.06–0.21	0.19–0.65
	Riceberry rice	0.002	0.005	0.007	0.016			0.12–0.26	0.36–0.79
STER	Jasmine rice	0.001	0.001	0.001	0.003	NA	0.1 (TDI) ^d	0.46–1.07	1.30–3.01
	Black sticky rice	0.001	0.001	0.006	0.007			1.01–1.56	5.99–6.83
	Riceberry rice	0.0003	0.001	0.001	0.002			0.32–0.60	0.97–1.82

^a LB, lower bound estimate calculated as the mean concentration for all samples; the undetected values and the unquantified values were replaced by zero and the LOD, respectively.

^b UB, upper bound estimate calculated as the mean concentration for all samples; the undetected values and the unquantified values were replaced by the LOD and the LOQ, respectively.

^c NA, not available.

^d Tolerable daily intake (TDI) with reference to TDI of HT-2 toxin of 0.1 $\mu\text{g}/\text{kg}$ of BW/day.

in all types of samples except brown rice, red rice, and barley. The prevalence of mycotoxin contamination differed among sample types. BEA, DAS, ZEA, and AFB₁ were found most frequently in all contaminated samples. However, the concentrations of regulated mycotoxins were below the established limits. The European Commission (7) has established its regulatory concentration for AFB₁ and total aflatoxins in cereals at 2 and 4 $\mu\text{g}/\text{kg}$, respectively. The maximum limit for ZEA in rice was set at 100 $\mu\text{g}/\text{kg}$ (6).

In this study, the EDI and the HQ were used to assess the health risk from rice and barley for adult Thai consumers. The EDIs of AFB₁, AFG₁, ZEA, BEA, DAS, and STER were below the PMTDI values established by JECFA. The risk assessment (HQ) value for ZEA residue was <100% in rice, which indicates no obvious risk to consumers in Thailand. For AFB₁, the risk assessment calculation was performed separately by determination of the population risk. AFB₁ contamination in GABA brown rice, black sticky rice, and riceberry rice was 1.38, 1.44, and 1.48 $\mu\text{g}/\text{kg}$, respectively, resulted in an estimated exposure of 4.14, 4.32, and 4.44 ng/kg of body weight per

day. Therefore, the estimated cancer risk to the Thai population through consumption of GABA brown rice, black sticky rice, and riceberry rice was 0.101, 0.105, and 0.109 cases of liver cancer per 100,000 persons per nanogram of AFB₁ per kilogram of body weight per day. Thus, only one liver cancer case per one million persons would be due to consumption of AFB₁ in rice. However, study of emerging mycotoxins in rice, such as BEA and STER, are warranted.

In summary, in this study the LC-MS/MS method was adequate for quantification of 16 mycotoxins in samples of rice and barley. AFB₁ and ZEA concentrations in the analyzed samples were lower than the maximum regulatory limits set by the European Commission. These results also suggest that the risk of harmful mycotoxin exposure via the consumption of barley and these nine kinds of rice (black sticky rice, brown rice, GABA rice, Japanese rice, jasmine rice, red rice, riceberry rice, white rice, and white sticky rice) is very low for Thai consumers. Further studies with a larger sample size are needed to confirm these data because

the samples may not be statistically representative of all types of rice.

ACKNOWLEDGMENTS

This work was partially supported by the Center for Advanced Studies for Agriculture and Food (Institute for Advanced Studies, Kasetsart University) under the Higher Education Research Promotion and National Research University Project of Thailand, Office of the Higher Education Commission, Ministry of Education, Thailand.

REFERENCES

- Arroyo-Manzanares, N., J. F. Huertas-Pérez, A. M. García-Campaña, and L. Gámiz-Gracia. 2014. Simple methodology for the determination of mycotoxins in pseudocereals, spelt and rice. *Food Control* 36:94–101.
- Bansal, J., P. Pantazopoulos, J. Tam, P. Caviovic, K. Kwong, and A. M. Turcotte. 2011. Surveys of rice sold in Canada for aflatoxins, ochratoxin A and fumonisins. *Food Addit. Contam.* 28:767–774.
- Bryden, W. L. 2012. Mycotoxin contamination of the feed supply chain: implications for animal productivity and feed security. *Anim. Feed Sci. Technol.* 173:134–158.
- Chinaphuti, A., and S. Aukkasarakul. 2009. Contamination of aflatoxin B₁ and ochratoxin A in brown rice and white rice. *Thai Rice Res. J.* 3:57–65. (In Thai.)
- European Commission. 2002. Commission Decision of 12 August 2002 implementing Council Directive 96/23/EC concerning the performance of analytical methods and the interpretation of results. 2002/657/EC. *Off. J. Eur. Communities* L 221:8–36.
- European Commission. 2006. Commission Regulation (EC) No 1881/2006 of 19 December 2006 setting maximum levels for certain contaminants in foodstuffs. *Off. J. Eur. Union* L 364:5–24.
- European Commission. 2010. Commission Regulation (EC) No 165/2010 of 26 February 2010 amending Regulation (EC) No 1881/2006 setting maximum levels for certain contaminants in foodstuffs as regards aflatoxins. *Off. J. Eur. Union* L 50:8–12.
- Food and Agriculture Organization of the United Nations. 2017. *Rice Market Monitor*, vol. 20, issue 4. Available at: <http://www.fao.org/economic/est/publications/rice-publications/rice-market-monitor-rmm/en/>. Accessed 15 May 2018.
- Institute of Nutrition, Mahidol University. 2015. Food consumption data of Thailand. Available at: <http://tinyurl.com/yc3ryrkl>. Accessed 8 May 2018. (In Thai.)
- Iqbal, S. Z., M. R. Asi, U. Hanif, M. Zuber, and S. Jinap. 2016. The presence of aflatoxins and ochratoxin A in rice and rice products; and evaluation of dietary intake. *Food Chem.* 210:135–140.
- Jettanajit, A., and T. Nhujak. 2016. Determination of mycotoxins in brown rice using QuEChERS sample preparation and UHPLC–MS–MS. *J. Chromatogr. Sci.* 54:720–729.
- Joint FAO/WHO Expert Committee on Food Additives. 1998. Safety evaluation of certain food additives and contaminants. Prepared by the forty-ninth meeting of the Joint FAO/WHO Expert Committee on Food Additives (JECFA). WHO Food Additives Series 40. World Health Organization, Geneva.
- Joint FAO/WHO Expert Committee on Food Additives. 2010. Summary and conclusions. Joint FAO/WHO Expert Committee on Food Additives seventy-second meeting, Rome, 16 to 25 February 2010. Food and Agriculture Committee of the United Nations, Rome.
- Kim, D.-H., S.-Y. Hong, J. W. Kang, S. M. Cho, K. R. Lee, T. K. An, C. Lee, and S. H. Chung. 2017. Simultaneous determination of multi-mycotoxins in cereal grains collected from South Korea by LC/MS/MS. *Toxins* 9:106. <https://doi.org/10.3390/toxins9030106>.
- Koesukwiwat, U., K. Sanguankaew, and N. Leepipatpiboon. 2014. Evaluation of a modified QuEChERS method for analysis of mycotoxins in rice. *Food Chem.* 153:44–51.
- Lim, C. W., T. Yoshinari, J. Layne, and S. H. Chan. 2015. Multi-mycotoxin screening reveals separate occurrence of aflatoxins and ochratoxin A in Asian rice. *J. Agric. Food Chem.* 63:3104–3113.
- Logrieco, A. F., J. D. Miller, M. Eskola, R. Krška, A. Ayalew, R. Bandyopadhyay, P. Battilani, D. Bhatnagar, S. Chulze, S. D. Saeger, P. Li, G. Perrone, A. Poapolathep, E. S. Rahayu, G. S. Shephard, F. Stepman, H. Zhang, and J. F. Leslie. 2018. The Mycotox Charter: increasing awareness of, and concerted action for, minimizing mycotoxin exposure worldwide. *Toxins* 10:149. <https://doi.org/10.3390/toxins10040149>.
- Miličević, D. R., M. Škrinjar, and T. Baltić. 2010. Real and perceived risks for mycotoxin contamination in foods and feeds: challenges for food safety control. *Toxins* 2:572–592.
- National Bureau of Agricultural Commodity and Food Standards. 2016. Food consumption data of Thailand. Bureau of Product Standards and Quality System, Bangkok, Thailand. Available at: <http://tinyurl.com/y8sqshus6>. Accessed 8 May 2018. (In Thai.)
- Nguyen, M. T., M. Tozlovanu, T. L. Tran, and A. Pфеhi-Leszkwicz. 2007. Occurrence of aflatoxin B₁, citrinin and ochratoxin A in rice in five provinces of the central region of Vietnam. *Food Chem.* 105:42–47.
- Panrapee, I., K. Phakpoom, M. Thanapoom, A. Nampeung, and M. Warapa. 2016. Exposure to aflatoxin B₁ in Thailand by consumption of brown and color rice. *Mycotoxin Res.* 32:19–25.
- Park, J. W., S. Y. Choi, H. J. Hwang, and Y. B. Kim. 2005. Fungal mycoflora and mycotoxins in Korean polished rice destined for humans. *Int. J. Food Microbiol.* 103:305–314.
- Puangkham, S., A. Poapolathep, U. Jermnak, K. Imsilp, P. Tanhan, C. Choekjaroenrat, and S. Poapolathep. 2017. Monitoring and health risk of mycotoxins in imported wines and beers consumed in Thailand. *World Mycotoxin J.* 10:401–409.
- Reddy, K. R. N., C. S. Reddy, H. K. Abbas, C. A. Abel, and K. Muralidharan. 2008. Mycotoxigenic fungi, mycotoxins, and management of rice grains. *Toxin Rev.* 27:287–317.
- Reddy, K. R. N., C. S. Reddy, and K. Muralidharan. 2009. Potential of botanicals and biocontrol agents on growth and aflatoxin production by *Aspergillus flavus* infecting rice grains. *Food Control* 20:173–178.
- Sales, A. C., and T. Yoshizawa. 2005. Updated profile of aflatoxin and *Aspergillus* section Flavi contamination in rice and its by-products from the Philippines. *Food Addit. Contam.* 22:429–426.
- Sun, X. D., P. Su, and H. Shan. 2017. Mycotoxin contamination of rice in China. *J. Food Sci.* 82:573–584.
- Suprasert, D. 2008. Contamination of aflatoxin in cereal grains. *Thai. J. Health Prom. Environ. Health* 31:102–107. (In Thai.)
- Tanaka, K., Y. Sago, Y. Zheng, H. Nakagawa, and M. Kushiro. 2007. Mycotoxins in rice. *Int. J. Food Microbiol.* 119:59–66.
- Toteja, G. S., A. Mukherjee, S. Diwakar, P. Singh, B. N. Saxena, K. K. Sinh, N. Kumar, K. V. Nagaraja, G. Bai, C. A. K. Prasad, S. Vanchinathan, R. Roy, and S. Sarkar. 2006. Aflatoxin B(1) contamination of parboiled rice samples collected from different states of India: a multi-centre study. *Food Addit. Contam.* 23:411–414.
- U.S. Environmental Protection Agency. 2011. Exposure factors handbook: 2011 edition. Office of Research and Development, National Center for Environmental Assessment, Washington, DC.
- Zimmer, I., E. Usleber, H. Klaffke, R. Weber, P. Majerus, H. Otteneder, and E. Märklbauer. 2008. Fumonisin intake of the German consumer. *Mycotoxin Res.* 24:40–52.